

PROPOSITION SUJETS DE THESES
CONTRATS DOCTORAUX 2020-2023

Appel ciblé (merci de cocher la case correspondante) :

- Contrat doctoral ministériel ED 536
 - Contrat doctoral ministériel ED 537
 - Contrat doctoral EUR Implantéus
-

Directeur de thèse : Edith GABRIEL, BioSP INRAE

Co-directeur éventuel : Dino IENCO, UMR TETIS

Co-encadrant éventuel : Eric VERDIN, PV INRAE

Titre en français : Une épidémio-surveillance augmentée de *Xylella fastidiosa* basée sur le risque

Titre en anglais : Enhanced risk-based surveillance of *Xylella fastidiosa*

Mots-clés :

Machine learning; Spatial statistics, Optimization algorithms, Epidemiology

Co tutelle : ~~Oui~~ Non **Pays :**

Opportunités de mobilité à l'international du doctorant dans le cadre de sa thèse : oui - non

Profil du candidat :

Strong understanding of machine learning;
Good coding skill in R (and/or Python, C++);
Knowledge on spatial statistics concepts would be a plus;
Interest for reading research papers;
Interest for applied mathematics and epidemiology

Présentation détaillée du sujet :

Martinetti and Soubeyrand (2018) proposed a risk-based surveillance strategy, based on a combination of machine learning techniques and network analysis, for understanding the main abiotic drivers of infections caused by *Xylella fastidiosa* and producing risk maps. Their predictors for explaining the presence of *Xylella fastidiosa* are mainly related to climate variables (precipitation seasonality, temperature in winter, solar radiation in summer...). Recent works, see e.g. Del Coco et al. (2020), reveal significant differences in soil and leaf ionome composition between safe areas and in presence of *Xylella fastidiosa*. Predictive models will be proposed on the basis of such new variables. Predictions will then be compared to the ones of Martinetti and Soubeyrand (2018). We will focus on proxies (eventually made of a combination of covariables) that allow large spatial coverage and cost reduction (in comparison with human inspections and specific biotic sensors).

Numerous factors potentially inform *Xylella fastidiosa* presence and dissemination (more than 100 factors, e.g. related to plant-health sanitary alerts, meteorological predictions, land use, satellite environmental data, abiotic sensor data...) and these factors are observed at various space-time scales and degrees of specificity. We thus need predictive models that deal with target variables and covariables that are potentially spatially and temporally autocorrelated and that can be noisy, more or less reliable, and collected at diverse spatial resolution x coverage x density. Machine learning, more specifically random forest approaches, has been successful in geoscience and spatial predictions, as well as in the study of systems that are dynamics. See e.g. Reichstein et al. (2019) and Salcedo-Sanz et al. (2020) for reviews on machine learning methods and information fusion in geoscience. However, machine learning methods rarely account for spatial and/or temporal dependencies, and hence high risk to lead to wrong extrapolation, interpretation, causal relation, and ignorance of confounding effects.

Several attempts to account for spatial or temporal dependencies have been made, but the mainstream approaches are based on hand-crafted features. In this project we want to go further, by jointly considering both the spatial and temporal dimensions to deal with spatio-temporal autocorrelation. Spatio-temporal dependencies will be overcome by deploying machine learning approaches (Meyer et al., 2019; Szatmári and Pásztor, 2019), including species distribution models (Norberg et al., 2019) and tailor them to the heterogeneity of the data and to our ES objectives: risk prediction and risk factor identification.

Coupling risk maps provided by our predictive models and optimization or allocation algorithms will then allow us to identify optimal risk-based surveillance (Mastin et al., 2020). However, instead of considering static situations, we will propose sequential sampling strategies conditional on the risk but also on previous samples, by mutually feeding risk mapping and data collection in a loop. Developing such a dynamic vision of sampling is especially paramount for *Xylella fastidiosa*: we know that a mild winter can favour bacteria expansion and symptom expression in regions beyond the usual suitability envelope, and surveys should be adapted accordingly.

Domaine / Thématique :

Machine learning ; statistics / Epidemiology

Objectif :

This project aims at finding new indicators and sampling strategies for ES across several horizons of space and time by (i) leveraging the heterogeneity and the mass of data, (ii) revealing links between potential predictors and observations of pathogens from heterogeneous observation strategies, and (iii) extrapolating and predicting the presence of a pathogen or the risk of its presence.

- We want to develop a generic supervised learning approach to conceive an ES informed by massive and heterogeneous data.
- Because designing efficient ES of invasive pathogens, as *Xylella fastidiosa*, relies on the knowledge of risk factors and their interactions, we will focus on finding new indicators that will be generated by the above-mentioned supervised learning approach.

Contexte et enjeux :

Invasive plant pathogens cause serious environmental and ecological damages with major economic impacts. *Xylella fastidiosa* is one of them, causing serious diseases in a wide range of plants such as grapevine and olive trees. Eradication management of *Xylella fastidiosa* currently involves surveying for early field symptoms, for plants that harbour the bacterium and then removing them and establishing demarcated areas.

Eliminating pesticides from plant protection involves a multitude of decisions for building and maintaining agro-food systems based on ecological and physical processes that favour plant health. The decisions required by farmers to deploy effectively prophylaxis are based on the availability of expert knowledge combined with numerous environmental variables whose states can be continuously updated via environmental monitoring – as part of systems referred to as epidemiological surveillance (ES).

In the case of olive tree, the Leccino cultivar has demonstrated tolerance to the disease (D’Attoma et al., 2019) – that seems to be associated with a specific micronutrient content of the plant and a specific microbial composition of its endophytes. Future surveillance could thus focus on endophytic composition of trees in regions with climates and soils most favourable for Olive Quick Decline symptom expression. This would give prophylaxis the needed time to alter plant nutritional states and endophytes to mimic those of the Leccino cultivar.

Surveys cover large and complex landscapes. The ES therefore integrates multiple sources of data that potentially inform the presence (or risk of presence) of the pathogen. Environmental variables monitored in ES typically include abiotic conditions that are mechanistically relevant to disease epidemiology (i.e., temperature, relative humidity, leaf wetness...) and traces of the pathogen (symptoms on local crops; changes in leaf ionome composition...).

Conceiving such an ES, incorporating a multitude of factors, across several horizons of space and time is thus challenging.

This PhD project will be supervised by two faculties working in two labs involved in IMPLANTEUS (BioSP and PV) and embodies one of the crucial features of the EUR: the interdisciplinarity. Moreover, the applied dimension of the project concerns a plant pathogen, which has been detected at several places throughout the Mediterranean basin in the past few years and which is a major threat for all the region from agricultural, social and economic perspectives. Finally, the expected results of the research that we propose related to the epidemiological surveillance of *Xylella fastidiosa* are paramount in the perspectives of large-scale disease management and public policy support.

Références bibliographiques :

D'Attoma G et al. (2019) Itonomic Differences between Susceptible and Resistant Olive Cultivars Infected by *Xylella fastidiosa* in the Outbreak Area of Salento, Italy. *Pathogens*, 8(4):272

Del Coco L et al. (2020) Soil and Leaf Ionome Heterogeneity in *Xylella fastidiosa* Subsp. Pauca-Infected, Non-Infected and Treated Olive Groves in Apulia, Italy. *Plants*, 9(6), 760.

Martinetti D and Soubeyrand S (2018) Identifying Lookouts for Epidemio-Surveillance: Application to the Emergence of *Xylella fastidiosa* in France. *Phytopathology*, 109(2):265–76.

Mastin J et al. (2020) Optimising risk-based surveillance for early detection of invasive plant pathogens. *PLOS Biology*, 18(10): e3000863.

Meyer H et al. (2019) Importance of spatial predictor variable selection in machine learning applications – Moving from data reproduction to spatial prediction. *Ecological Modelling*. 411:108815.

Norberg A et al. (2019) A comprehensive evaluation of predictive performance of 33 species distribution models at species and community levels. *Ecological Monographs*, 89(3):e01370.

Reichstein M et al. (2019) Deep learning and process understanding for data-driven Earth system science. *Nature*, 566(7743), 195–204.

Salcedo-Sanz S et al. (2020) Machine learning information fusion in Earth observation: A comprehensive review of methods, applications and data sources. *Information Fusion*, 63, 256–272.

Szatmári G and Pásztor L. (2019) Comparison of various uncertainty modelling approaches based on geostatistics and machine learning algorithms. *Geoderma*. 337, 1329–1340.

Les sujets devront être adressés à

gestion-ed@univ-avignon.fr

Avant le 6 avril 2020